

## POLYMORPHONUCLEAR LEUKOCYTE RHEOLOGY, CYTOSOLIC $Ca^{2+}$ CONTENT, BETA<sub>2</sub>-INTEGRIN EXPRESSION AND OXIDATIVE STRESS IN HYPERTENSION

Gregorio Caimi, Eugenia Hopps, Rosalia Lo Presti

### INTRODUCTION

An elevated leukocyte count is associated to arterial hypertension, but in this brief review we focus our attention on some functional aspects of polymorphonuclear leukocytes (PMNs), considering in particular our data previously published on this topic. Leukocytes have a likely role in the pathogenesis of organ injury accompanying arterial hypertension. Monocytes and neutrophils can in fact participate in the progression of vascular lesions and in the pathophysiology of their ischaemic complications.

### LEUKOCYTE RHEOLOGY

Hypertensives have a whole-blood hyperviscosity, due to increased leukocyte count, erythrocyte rigidity and hematocrit, and also to a raise of plasma viscosity, fibrinogen concentration, erythrocyte and platelet ag-

gregation. As blood viscosity influences peripheral vascular resistances, changes of the haemorheological parameters may contribute to the development of hypertension and can facilitate myocardial and vascular remodelling<sup>1</sup>.

In hypertensives the reduced membrane fluidity of blood cells affects carrier activity: in PMNs alterations of sodium and calcium plasma membrane pumps have been observed. Higher sodium cytosolic concentrations promote leukocyte activation, superoxide anion synthesis, adhesion to endothelium and progressive organ injury<sup>2</sup>. Several researches have previously demonstrated normal cytosolic  $Ca^{2+}$  concentrations in hypertensive monocytes and PMNs<sup>3-5</sup>. On the contrary, we observed increased cytosolic  $Ca^{2+}$  concentrations in PMNs<sup>6</sup>, as well as in platelets and erythrocytes<sup>7</sup>. This datum was confirmed by more recent studies in insulin-resistant hypertensive men<sup>8</sup>.

Dipartimento di Medicina Interna, Malattie Cardiovascolari e Nefrourologiche  
Cattedra di Semeiotica e Metodologia Medica  
Università di Palermo, Italy

#### Corresponding author:

Prof. Gregorio Caimi  
Via Leonardo da Vinci, 52  
90145 Palermo, Italy  
Fax: +39 91 6554535; e-mail: caimigre@unipa.it

In our research, at baseline, no correlation was present between PMN membrane fluidity and cytosolic  $\text{Ca}^{2+}$  content or between membrane fluidity and arterial pressure in hypertensives<sup>7,9</sup>, whereas PMN membrane fluidity was decreased in hypertensives compared to normals<sup>10</sup>. After *in vitro* activation with PMA the initial relative flow rate (obtained by filtering unfractionated WBC and PMN suspension using a St. George's Filtrometer) was decreased in normals and hypertensives, but no other modifications were found. After activation with fMLP,  $\text{Ca}^{2+}$  content was furtherly increased only in hypertensives<sup>11,12</sup>.

### LEUKOCYTE ACTIVATION

In hypertensives, leukocytes have higher activation levels than in normal subjects<sup>13</sup>. Stimulated granulocytes release reactive oxygen species (ROS) which may inactivate nitric oxide (NO), arachidonic acid metabolites which induce vascular inflammation, and cathepsin G which stimulates the angiotensin system<sup>14</sup>. Primed PMNs may also initiate platelet aggregation<sup>14</sup> and provoke leukocyte adhesion to the endothelium promoting the hyperexpression of adhesion molecules<sup>15</sup>, and the NO inactivation by superoxide anion. NO is in fact an endogenous inhibitor of leukocytes adhesion in postcapillary venules<sup>16</sup>.

Subjects with essential hypertension have increased plasma levels of leukocyte elastase<sup>17,18</sup>, that is considered a sensitive diagnostic marker of cardiovascular disease. Higher elastase levels are associated with the presence of atheromatous plaques<sup>19</sup>

and are related to arterial stiffness<sup>18</sup>. In spontaneously hypertensive rats (SHRs) circulating PMNs have lower alkaline phosphatase content, while the plasma levels of this enzyme are elevated in these models<sup>16,20</sup>; this finding is another indicator of leukocyte activation and spontaneous degranulation and reflects the alteration of leukocyte function too.

### LEUKOCYTES AND OXIDATIVE STRESS

Leukocytes activated by soluble or cellular stimuli start the respiratory burst, characterized by increased oxygen consumption and generation of superoxide ( $\text{O}_2^-$ ), hydrogen peroxide ( $\text{H}_2\text{O}_2$ ) and other ROS.  $\text{O}_2^-$  synthesis by hypertensive PMNs is related to blood pressure [21]. ROS contribute to hypertension and its complications causing endothelial damage, oxidizing circulating low density lipoproteins and degrading NO. Serum NO levels and arterial pressure are negatively correlated<sup>22</sup>. The decreased availability of NO has negative effects on vascular tone and influences platelet aggregation, leukocyte adhesion and expression of adhesion molecules<sup>23</sup>. Oxidative stress in PMNs is related to arterial mean pressure, as well as oxidative stress in monocytes is related to C-reactive protein (CRP)<sup>24</sup>.

A research regarding Sabra hypertension-resistant rats (SBN) and hypertension-prone rats (SBH) has shown that in pre-hypertensive SBH oxidative stress and primed PMNs are present. After salt-loading for four weeks, SBH rats developed hypertension and their PMNs pro-

duce elevated levels of ROS<sup>25</sup>. A long-term antioxidant treatment with Tempol (a membrane-permeable SOD mimetic), initiated in pre-hypertensive SHR, can attenuate vascular oxidative stress and prevent the age-related elevation of blood pressure in this animal model of genetic hypertension<sup>26</sup>.

A recent study revealed a spontaneous up-regulation of CD11a and CD18 in circulating PMNs and an increased ROS generation by T-lymphocytes and monocytes in the peripheral blood of obese Zucker rats, before the development of hypertension<sup>27</sup>. This data suggest that primed PMNs, oxidative stress and inflammation anticipate and possibly contribute to the development of hypertension and its complications.

In Sabra rats salt-loaded for sixty days, the increase of blood pressure can be inhibited by phenylarsine oxide, an NADPH oxidase inhibitor. This finding suggests that primed PMNs play a role in the development of hypertension via activation of NADPH oxidase and production of ROS<sup>28</sup>. Because the leukocyte NADPH oxidase is localized in the plasma membrane, the alterations of its activity may be related to the membrane abnormalities observed in hypertension<sup>29</sup>.

## LEUKOCYTE INTEGRINS

In hypertensives, the increased expression of adhesion molecules on endothelial cells and leukocytes promotes leukocyte adhesion to the endothelium<sup>30</sup>. A research on SHR has demonstrated a decreased endothelial expression of P-selectin<sup>31</sup>. On the

contrary, another study showed a venular P-selectin hyperexpression that is considered responsible for an increased rolling of PMNs<sup>32</sup>. The increased level of soluble E- and P-selectin in malignant and renovascular hypertension is due to platelet activation<sup>33</sup>. E-selectin, an indicator of organ injury, is related to blood pressure even in hypertensives without complications<sup>34</sup>. Benidipine, a calcium channel blocker, decreases arterial pressure and the concentrations of E- and P-selectin<sup>35</sup>. In coronary endothelial cells derived from human heart, angiotensin II induces a concentration-dependent increase in E-selectin expression via activation of AT1 receptors. So, the E-selectin-mediated leukocyte adhesion to the endothelium can be blocked using AT1 receptor antagonists<sup>36</sup>. The infusion of angiotensin II *in vivo* induces high levels of soluble ICAM-1 (Intracellular Adhesion Molecules-1) in hypertensives and normals<sup>37</sup>. Hypertension increases cytokine expression and concentrations of ICAM-1 and MCP-1 (monocyte chemoattractant peptide-1), responsible for the inflammatory cell infiltration of the subendothelium<sup>38</sup>: exposing mouse carotid arteries at high intraluminal pressure, after 24 hours an increased mRNA expression of MCP-1, IL-6, chemokines and VCAM-1 (Vascular Cell Adhesion Molecule-1) is detectable in monocytes<sup>39</sup>. VCAM-1 is positively related to systolic blood pressure<sup>40</sup>. The induction of hypertension in genetically normal rats can lead to an overexpression of the CD18 integrin on circulating helper T-cells<sup>41</sup>. In obese hypertensives an increased expression of CD11b on circulating monocytes and of CD68 on macrophages

in the adipose tissue was found<sup>42</sup>. Studies evaluating PMN  $\beta_2$ -integrins showed at baseline a phenotypical hyperexpression of CD11a, CD11b and CD18, but not of CD11c. After *in vitro* activation an increase of CD11b, CD11c and CD18 has been demonstrated, but also a decrease of CD11a, especially with PMA<sup>43</sup>. The integrin overexpression on PMNs at baseline can be due to the spontaneous granulocyte activation. The fall in CD11a expression after activation might be due to different mechanisms: because CD11a is constitutively expressed on PMN surface and it is not internalized during activation, its decrease may be related to an altered cleavage or to a dysregulated phosphorylation/dephosphorylation process<sup>44</sup>. However, up to now the mechanisms that explain the integrin pattern are only partially known.

## REFERENCES

- Shliakhto EV, Moiseeva OM, Liasnikova EA, Villeval'de SV, Emel'ianov IV. Rheological properties of blood and endothelial function in patients with hypertensive disease. *Kardiologija* 2004; 44:20-23.
- Schmid-Schonbein GW, Seiffge D, DeLano FA, Shen K, Zweifach BW. Leukocyte counts and activation in Spontaneously Hypertensive and Normotensive Rats. *Hypertension* 1991; 17:323-330.
- Lew PD, Favre L, Waldvogel FA, Vallotton MB. Cytosolic free calcium and intracellular calcium stores in neutrophils from hypertensive subjects. *Clin Sci* 1985; 69:227-230.
- Shore AC, Beynon GW, Jones JC, Markandu ND, Sagnella GA, MacGregor GA. Mononuclear leukocyte intracellular free calcium – Does it correlate with blood pressure? *J Hypertens* 1985; 3:183-187.
- Bing RF, Heagerty AM, Jackson JA, Thurston H, Swales JD. Leukocyte ionized calcium and sodium content and blood pressure in humans. *Hypertension* 1986; 8:483-488.
- Caimi G, Canino B, Montana M, Ventimiglia G, Catania A, Lo Presti R. Polymorphonuclear leukocyte membrane fluidity and cytosolic Ca<sup>2+</sup> content in different clinical conditions. *Clin Hemorheol Microcirc* 1997; 17:217-223.
- Caimi G, Contomo A, Serra A, Catania A, Lo Presti R, Sarno A, Cerasola G. Red cell metabolic parameters and rheological determinants in essential hypertension. *Clin Hemorheol* 1993; 13:35-44.
- Sela S, Shurtz-Swirski R, Farah R, Levy R, Shapiro G, Chezari J, Shasha SM, Kristal B. A link between polymorphonuclear leukocyte intracellular calcium, plasma insulin and essential hypertension. *Am J Hypertens* 2002; 15:291-295.
- Caimi G. Erythrocyte, platelet and polymorphonuclear leukocyte membrane dynamic properties in essential hypertension. *Clin Hemorheol Microcirc* 1997; 17:199-208.
- Lo Presti R, Carollo C, Canino B, Montana M, Romano A, Catania A, Caimi G. Polymorphonuclear leukocyte membrane fluidity and cytosolic Ca<sup>2+</sup> content at baseline and after activation in essential hypertension. *Trace Elem Electrolytes* 2005; 22:207-210.
- Caimi G, Lo Presti R, Canino B, Montana M, Ferrara L, Oddo G, Vetimiglia G, Cerasola G. Essential hypertension: leukocyte rheology and polymorphonuclear cytosolic Ca<sup>2+</sup> content at baseline and after activation. *Clin Hemorheol Microcirc* 1998; 19:281-289.
- Caimi G, Lo Presti R, Canino B, Montana M, Cerasola G. Essential hypertension: polymorphonuclear leukocyte membrane fluidity at baseline and after chemotactic activation. *Am J Hypertens* 1999; 12:947-948.
- Chatterjee M, Saluja R, Kanneganti S, Chinta S, Dikshit M. Biochemical and molecular evaluation of neutrophil NOS in spontaneously hypertensive rats. *Cell Mol Biol* 2007; 15:84-93.
- Mugge A, Lopez JAG. Do leukocytes have a role in hypertension? *Hypertension* 1991; 17:331-333.
- Pardon NJ, Wilkinson R, Thomas TH. Rapid fusion of granules with neutrophil cell membranes in hypertensive patients may increase vascular damage. *Am J Hypertens* 2001; 14:927-933.
- Shen K, Sung KL, Whittemore, DeLano FA, Zweifach BW, Schmid-Schonbein GW. Properties of circulating leukocytes in spontaneously hypertensive rats. *Biochem Cell Biol* 1995; 73:491-500.
- Jackson MH, Collier A, Nicoll JJ, Muir AL, Dawes J, Clarke BF, Bell D. Neutrophil count and activation in vascular disease. *Scott Med J* 1992; 37:41-43.
- Yasmin SW, McEniery CM, Dakham Z, Pusalkar P, Maki-Petaja K, Ashby MJ, Cockcroft JR, Wilkinson IB. Matrix metalloproteinase-9 (MMP-9), MMP-2, and serum elastase activity are associated with systolic hypertension and arterial stiffness. *Arterioscler Thromb Vasc Biol* 2005; 25:372-378.
- Amaro A, Gude F, Gonzalez-Juanatery R, Iglesias C, Fernandez-Vasquez F, Garcia-Acuna J, Gil M. Plasma leukocyte elastase concentration in angiographically diagnosed coronary artery disease. *Eur Heart J* 1995; 16:615-622.
- Kristal B, Shurtz-Swirski R, Chezari J, Manaster J, Levy R, Shapiro G, Weissman I, Shasha SM, Sela S. Participation of peripheral polymorphonuclear leukocytes in the oxidative stress and inflammation in patients with essential hypertension. *Am J Hypertens* 1998; 11:921-928.
- Mazor R, Shurtz-Swirski R, Sela S, Shapiro G, Kristal B. Polymorphonuclear leukocyte priming

- and counts are in correlation with blood pressure parameters. *Harefuah* 2006; 145:900-903.
22. Armas-Padilla MC, Armas-Hernández MJ, Sosa-Canache B, Cammarata R, Pacheco B, Guerrero J, Carvajal AR, Hernández-Hernández R, Israili ZH, Valasco M. Nitric oxide and malondialdehyde in human hypertension. *Am J Ther* 2007; 14:172-176.
  23. Portaluppi F, Boari B, Manfredini R. Oxidative stress in essential hypertension. *Curr Pharm Des* 2004; 10:1695-1698.
  24. Yasunari K, Maeda K, Nakamura M, Yoshikawa J. Oxidative stress in leukocyte is a possible link between blood pressure, blood glucose and C-reactive protein. *Hypertension* 2002; 39:777-780.
  25. Sela S, Mazor R, Amsalam M, Yagil C, Yagil Y, Kristal B. Primed polymorphonuclear leukocytes, oxidative stress, and inflammation antecede hypertension in the Sabra rat. *Hypertension* 2004; 44:764-769.
  26. Nabha L, Garbern JC, Buller CL, Charpie JR. Vascular oxidative stress precedes high blood pressure in spontaneously hypertensive rats. *Clinical and Experimental Hypertension* 2005; 27:71-82.
  27. Kim CH, Vaziri ND, Rodriguez-Iturbe B. Integrin expression and H<sub>2</sub>O<sub>2</sub> production in circulating and splenic leukocytes of obese rats. *Obesity* 2007; 15:2209-2216.
  28. Mazor R, Kristal B, Cohen-Mazor M, Yagil C, Yagil Y, Sela S. The polymorphonuclear leukocyte contributes to the development of hypertension in the Sabra rat. *J Hypertens* 2007; 25: 2249-2256.
  29. Sagar S, Kallo IJ, Kaul N, Ganguly NK, Sharma BK. Oxygen free radicals in essential hypertension. *Mol Cell Biochem* 1992; 111:103-108.
  30. Buemi M, Allegra A, Aloisi C, Corica F, Alonci A, Ruello A, Montalto G, Frisina N. Cold pressor test raises serum concentrations of ICAM-1, VCAM-1 and E-selectin in normotensive and hypertensive patients. *Hypertension* 1997; 30:845-847.
  31. Suzuki H, Zweifach BJ, Forrest MJ, Schmid-Schonbein GW. Modification of leukocyte adhesion in spontaneously hypertensive rats by adrenal corticosteroids. *J Leuk Biol* 1995; 57: 20-27.
  32. Suematsu M, Suzuki H, Tamatani T, Iigou Y, DeLano FA, Miysaka M, Forrest MJ, Kannagi R, Zweifach BJ, Ishimura Y, Schmid-Schonbein GW. Impairment of Selectin-mediated leukocyte adhesion to venular endothelium in spontaneously hypertensive rats. *J Clin Invest* 1995; 96:2009-2016.
  33. Verhaar MC, Beutier JJ, Gaillard CA, Koomans HA, Fijnheer R, Rabelink TJ. Progressive vascular damage in hypertension is associated with increased levels of circulating P-Selectin. *J Hypertens* 1998; 16:45-50.
  34. Miller MA, Kerry SM, Cook DG, Cappuccio FP. Cellular adhesion molecules and blood pressure: interaction with sex in a multi-ethnic population. *J Hypertens* 2004; 22:705-711.
  35. Sanada H, Midorikawa S, Yatabe J, Sasaki Yatabe M, Katoh T, Baba T, Hashimoto S, Watanabe T. Elevation of serum soluble E- P-selectin in patients with hypertension is reversed by benidipine, a long-acting calcium channel blocker. *Hypertens Res* 2005; 28:871-878.
  36. Grafe M, Auch-Schwelk W, Zakrewicz A, Regitz-Zagrosek V, Bartsch P, Graf K, Loebe M, Gaethgens P, Fleck E. Angiotensin II-induced leukocyte adhesion on human coronary endothelial cells is mediated by E-selectin. *Circ Res* 1997; 81:801-811.
  37. Pastore L, Tessitore A, Martinotti S, Toniato E, Alesse E, Bravi MC, Ferri G, Desideri A, Gulino A, Santucci A. Angiotensin II stimulates intercellular adhesion molecule-1 (ICAM-1) expression by human vascular endothelial cells and increases soluble ICAM-1 release in vivo. *Circulation* 1990; 100:1646-1652.
  38. Madej A, Okopien B, Kowalski J, Haberk M, Herman ZS. Plasma concentrations of adhesion molecules and chemokines in patients with essential hypertension. *Pharmacol Rep* 2005; 57:878-881.
  39. Riou S, Mees B, Esposito B, Merval R, Vilar J, Stengel D, Ninio E, van Haperen R, de Crom R, Tedgui A, Lehoux S. High pressure promotes monocyte adhesion to the vascular wall. *Circ Res* 2007; 100:1226-1233.
  40. De Souza CA, Dengel DR, Macko RF, Cox K, Seals DR. Elevated levels of circulating cell adhesion molecules in uncomplicated essential hypertension. *Am J Hypertens* 1997; 10:1335-1341.
  41. Kim CH, Vaziri ND. Hypertension promotes integrin expression and reactive oxygen species generation by circulating leukocyte. *Kidney Int* 2005; 67:1462-1470.
  42. Boschmann M, Engeli S, Adams F, Gorzelniak K, Franke G, Klaua S, Kreuzberg U, Luedtke S, Ketzritz R, Sharma AM, Luft FC, Jordan J. Adipose tissue metabolism and CD11b expression on monocytes in obese hypertensives. *Hypertension* 2005; 46:130-136.
  43. Caimi G, Lo Presti R, Carollo C, Musso M, Porretto F, Canino B, Catania A, Cerasola G. Polymorphonuclear integrins, membrane fluidity and cytosolic Ca<sup>2+</sup> content after activation in essential hypertension. *Hypertension* 2000; 36:813-817.
  44. Caimi G. Polymorphonuclear leukocyte integrin profile in essential hypertension. *Am J Hypertens* 2000; 13:1051-1052.